ELSEVIER J. Biochem. Biophys. Methods 35 (1997) 209–210

JOURNAL OF biochemical and biophysical methods

Author Index

Alonso, S., see Romero, E.L. (35) 129

Cárdenas, G., see Escobar, J. (35) 57
Ceolín, M., Martínez, J.A. and Ermácora, M.R.
Time differential perturbed angular correlation of ^{111}In-EDTA linked to the single free cysteine of bovine serum albumin (35) 135
Chan, S.D.H., see Dill, K. (35) 197
Chatterji, D., see Nagesh, N. (35) 61
Chen, N. and Chrambach, A.
Capillary electrophoresis of DNA fragments in 9 to 20% uncrosslinked polyacrylamide gels: unique separating capacity hypothetically related to maintenance of random-coil DNA conformation independently of gel concentration (35) 175
Chrambach, A., see Chen, N. (35) 175
Cole, J.W., Hendler, R.W., Smith, P.D., Fredrickson, H.A., Pohida, T.J. and Friauf, W.S.
A high speed optical multichannel analyzer (35) 161
Córdoba, O.L., Linskens, S.B., Dacci, E. and Santomé, J.A.
'In gel' cleavage with cyanogen bromide for protein internal sequencing (35) 1

Dacci, E., see Córdoba, O.L. (35) 1
Dill, K., Chan, S.D.H. and Gibbs, T.W.
Detection of plasmids using DNA and RNA probes and the light-addressable potentiometric sensor (35) 197

Ermácora, M.R., see Ceolín, M. (35) 135
Escobar, J., Cárdenas, G. and Lissi, E.
Evaluation of the total content of lipid-soluble antioxidants in blood plasma samples employing a simple chemiluminescence quenching procedure (35) 57
Escrich, E., see Solanas, M. (35) 153

Fredrickson, H.A., see Cole, J.W. (35) 161
Friauf, W.S., see Cole, J.W. (35) 161
Fromm, M., see Gitter, A.H. (35) 81

Gemeiner, P., see Mislovičová, D. (35) 37
Gibbs, T.W., see Dill, K. (35) 197
Gill, D., see Uzan, B. (35) 49
Gitter, A.H., Schulzke, J.-D., Sorgenfrei, D. and Fromm, M.
Ussing chamber for high-frequency transmural impedance analysis of epithelial tissues (35) 81
Glazachev, Yu.I., see Khramtsov, V.V. (35) 115
Green, N.G., Morgan, H. and Milner, J.J.
Manipulation and trapping of sub-micron bioparticles using dielectrophoresis (35) 89

Hendler, R.W., see Cole, J.W. (35) 161
Holmsen, H., see Jensen, B.O. (35) 185

Ishibashi, O.
A new method to synthesize competitor RNAs for accurate analyses by competitive RT-PCR (35) 203

Jensen, B.O., Skeidsvoll, J. and Holmsen, H.
A polarographic method for measuring dissolved nitric oxide (35) 185
Johansson, P. and Wadsö, I.
A photo microcalorimetric system for studies of plant tissue (35) 103
Jurinke, C., see van den Boom, D. (35) 69

Kam, W.K., see Li, Q.-T. (35) 11
Khramtsov, V.V., Yelinova, V.I., Glazachev, Yu.I., Reznikov, V.A. and Zimmer, G.
Quantitative determination and reversible modification of thiols using imidazolidine biradical disulfide label (35) 115
Köster, H., see van den Boom, D. (35) 69
Kujau, M.J., Siebert, A. and Wölfl, S.

S0165-022X(97)00048-1

Design of leader sequences that improve the efficiency of the enzymatic synthesis of 2′-amino-pyrimidine RNA for in vitro selection (35) 141

Li, Q.-T. and Kam, W.K.
Steady-state fluorescence quenching for detecting acyl chain interdigitation in phosphatidylcholine vesicles (35) 11
Linskens, S.B., see Córdoba, O.L. (35) 1
Lissi, E., see Escobar, J. (35) 57

Makrigiorgos, G.M.
Detection of lipid peroxidation on erythrocytes using the excimer-forming property of a lipophilic BODIPY fluorescent dye (35) 23
Martínez, J.A., see Ceolín, M. (35) 135
Milner, J.J., see Green, N.G. (35) 89
Mislovičová, D., Vikartovská, A. and Gemeiner, P.
The glycosylated enzyme-binding assay for the study of the interaction of free and immobilized lectins with carbohydrates (35) 37
Morgan, H., see Green, N.G. (35) 89

Nagesh, N. and Chatterji, D.
A useful method for the synthesis of 8 azido-guanosine tri- and tetraphosphate: important substrates for many enzymatic reactions (35) 61

Pardo, M.F., see Romero, E.L. (35) 129
Pines, E., see Uzan, B. (35) 49
Pohida, T.J., see Cole, J.W. (35) 161
Porro, S., see Romero, E.L. (35) 129

Reznikov, V.A., see Khramtsov, V.V. (35) 115
Romero, E.L., Pardo, M.F., Porro, S. and Alonso, S.
Sialic acid measurement by a modified Aminoff method: a time-saving reduction in 2-thiobarbituric acid concentration (35) 129
Ruppert, A., see van den Boom, D. (35) 69

Santomé, J.A., see Córdoba, O.L. (35) 1
Schulzke, J.-D., see Gitter, A.H. (35) 81
Siebert, A., see Kujau, M.J. (35) 141
Skeidsvoll, J., see Jensen, B.O. (35) 185
Smith, P.D., see Cole, J.W. (35) 161
Solanas, M. and Escrich, E.
An improved protocol to increase sensitivity of Southern blot using dig-labelled DNA probes (35) 153
Sorgenfrei, D., see Gitter, A.H. (35) 81

Uzan, B., Pines, E. and Gill, D.
Removal of dissolved O_2 by a fluid membrane of Fomblin Y (perfluoro polyether): fluorescence of diphenylhexatriene in Fomblin Y (35) 49

van den Boom, D., Ruppert, A., Jurinke, C. and Köster, H.
Combined amplification and sequencing in a single reaction using two DNA polymerases with differential incorporation rates for dideoxynucleotides (35) 69
Vikartovská, A., see Mislovičová, D. (35) 37

Wadsö, I., see Johansson, P. (35) 103
Wölfl, S., see Kujau, M.J. (35) 141

Yelinova, V.I., see Khramtsov, V.V. (35) 115

Zimmer, G., see Khramtsov, V.V. (35) 115

ELSEVIER J. Biochem. Biophys. Methods 35 (1997) 211–212

JOURNAL OF biochemical and biophysical methods

Subject Index

Alternating current impedance, (2) 81
2′-amino-pyrimidine nucleotides, (3) 141

Bacteriorhodopsin, (3) 161
Bead cellulose, (1) 37
Bio-particle manipulation, (2) 89
Blood plasma, (1) 57, (2) 115
BODIPY-FL, (1) 23

Capillary electrophoresis, (3) 175, 185
Chemiluminescence, (1) 57, (3) 153
Clark electrode, (1) 49
Competitive RT-PCR, (3) 203
Concanavalin A, (1) 37
Cytochrome oxidase, (3) 161
C_{16}-BODIPY, (1) 23

Deoxygenation, (1) 49
Dielectrophoresis, (2) 89
Digoxigenin, (3) 153
Diphenylhexatriene, (1) 49
Disulfide biradical, (2) 115
DNA conformation, (3) 175
DNA fragments, (3) 175
DNA polymerases, (2) 69
DNA probes, (2) 197
DNA sequencing, (2) 69
Dye terminator, (2) 69

Edman degradation, (1) 1
Electroblotting, (1) 1
Endothelium-derived relaxing factor, (3) 185
Epithelial resistance, (2) 81
EPR spectroscopy, (2) 115
Erythrocytes, (1) 23, (2) 115

Flow microcalorimetry, (1) 37
Fluid membrane, (1) 49
Fluorescence quenching, (1) 11
Fomblin Y, (1) 49
Free radicals, (1) 57

Gel electrophoresis, (1) 1
Glutathione, (2) 115

High accuracy, (2) 135
HPLC, (2) 129
HT-29 cells, (2) 81
Human serum albumin, (2) 115
Hybridization, (2) 197
Hyperfine techniques, (2) 135

Imidazolidine nitroxide, (2) 115
'In gel' CNBr cleavage, (1) 1
In vitro selection, (3) 141
Inhibition assay, (1) 37
Interdigitation, (1) 11

Latex spheres, (2) 89
Lectin-glycosylated enzyme interaction, (1) 37
Light addressable potentiometric sensor (LAPS), (2) 197
Lipid-soluble, (1) 57
Lipophilic analogues, (1) 23
Long accurate PCR, (3) 203

Micro-electrodes, (2) 89
Mild chemical modification, (2) 135

NBD-PC, (1) 11
Neu-5-Ac, (2) 129
Nitric oxide, (3) 185
Non-radioactive, (3) 153
N_3-GTP, (1) 61
N_3-ppGpp, (1) 61

Optical multichannel analyzer, (3) 161
Oxidation, (3) 185
Oximetry, (3) 185

Perfluorocarbon, (1) 49
Phenyl boronate column, (1) 61
Picomole, (2) 129

S0165-022X(97)00049-3

Plasmid detection, (2) 197
Platelet aggregation, (3) 185
Polarography, (3) 185
Polymerase chain reaction (PCR), (2) 69
Primer design, (3) 141
Protein sequence analysis, (1) 1
Purification, (1) 61

Quantitative analyses, (3) 203

Reaction mechanism, (3) 161
Reverse transcription, (3) 141, 203
RNA probes, (2) 197
RNA stability, (3) 141
Rotational correlation times, (2) 135

Sensitivity, (3) 153
Separating capacity, (3) 175
Southern Blot, (3) 153
Subepithelial resistance, (2) 81
Synthesis, (1) 61

TDPAC, (2) 135
2-Thiobarbituric acid (TBA), (2) 129
Thiols, (2) 115
Tobacco mosaic virus, (2) 89
Transcription, (3) 141

Uncrosslinked polyacrylamide, (3) 175
Ussing chamber, (2) 81